

**BURLINGTON COUNTY ENDOSCOPY CENTER**  
**140 Mt. Holly Bypass, Unit 5**  
**Lumberton, NJ 08048**

**COLONOSCOPY INSTRUCTIONS – Evening/Morning Prep (Split Dose)**  
**CLENPIQ – PRESCRIPTION REQUIRED**

Date of Procedure: \_\_\_\_\_

Report Time for Procedure: \_\_\_\_\_

\_\_\_\_\_ E-Prescribed to Pharmacy

\_\_\_\_\_ Prescription given to patient

**If you DO NOT follow these instructions your procedure may be cancelled**

If you have any **questions**, or **you are unable to complete most of the prep given or have not moved your bowels**, please call the GCSJ office ***IMMEDIATELY*** at **609-265-1700**. If it is after 5 pm you will receive a callback from the Doctor On-Call for the evening.

**PRIOR TO YOUR PROCEDURE**: Be sure to get your Clenpiq Prep Kit at least “1” week before your procedure. **If you take ANY BLOOD THINNERS OR MEDICATION FOR DIABETES, follow the instructions given to you by the provider at your appointment OR by the nurse during your phone interview.**

**7 DAYS PRIOR TO THE PROCEDURE STOP THE FOLLOWING**: (Unless otherwise instructed by your physician)

**Fish Oil** , IRON or Multivitamin with iron, OTC Supplements

**THE ENTIRE DAY BEFORE THE PROCEDURE – (YOUR PREP DAY)**

You are to be on a CLEAR LIQUID DIET. **NO SOLID FOOD (NO RED or PURPLE COLORED PRODUCTS – ORANGE IS OK)**

This includes **only** the following:

1. Water
2. Clear Broth: chicken, beef, vegetable
3. Clear juices: apple, white grape, white cranberry
4. Soft drinks: soda, Gatorade, lemonade- **NO RED OR PURPLE**
5. Coffee or tea: (hot or iced) – **NO CREAM, MILK, or POWDERED CREAMER**
6. Jell-O- **NO RED OR PURPLE**
7. Popsicles (without fruit or cream), lemon sorbet, Italian Ice

**START PREP: Plan on being at home during your prep. For your comfort you may use Vaseline or moistened baby wipes around the anal area.**

**Begin** the prep after work **but no later** than 7:00 pm (day prior to the procedure)

1. Prep is already prepared **NO MIXING REQUIRED. DO NOT ADD ANY LIQUID TO THE PREP.** You must drink the entire contents of the container.
2. **Drink FIVE (5)**, eight ounce glasses of **ANY** clear liquid on the list. Drink this over the next 2 hours. You will most likely start having loose bowel movements within 1-2 hours after drinking the bowel prep and clear liquids. This may last for several hours.

Try to be up and about (active) as much as possible while drinking the solution. You may continue to drink clear liquids from the list during the prep time. It is important for you to prevent yourself from becoming dehydrated so KEEP DRINKING THE RECOMMENDED CLEAR LIQUIDS up until midnight.

**NO SMOKING CIGARETTES OR SMOKING MARIJUANA, INGESTING OF MARIJUANA, MEDICINAL MARIJUANA OR EDIBLES. NO HARD CANDY, LIFESAVERS, MINTS, OR GUM\*\*\*\*\***

**MORNING OF THE PROCEDURE : Your morning prep start will be 6 HOURS PRIOR TO YOUR PROCEDURE TIME. GCSJ will advise you when to begin the final dose of clenpiq.**

**\*\*\*\*\*You must finish your prep and not drink anything else within 5 hours of your procedure\*\*\*\*\***

**Begin** your final dose of Clenpiq at \_\_\_\_\_ a.m. on the morning of your procedure.

DRINK THE SECOND DOSE using the instructions from the previous evening. The instructions are exactly the same. Once you have consumed the prep, you should **immediately** follow it by drinking a quantity of **3 to 5**, 8 ounce glasses of any clear liquid over 1 hour.

You **MAY** take your morning medications after your last dose of the morning prep with a sip of water. **MEDS SHOULD BE TAKEN FOUR HOURS PRIOR TO PX.** However, if you are DIABETIC, you **MAY NOT** take your diabetes medication.

**You are required to have a responsible party over the age of 18 accompany you to provide transportation after your procedure. For the safety of our patients, we do not allow our patients to drive themselves home. We recommend that your driver wait for you during the visit to the surgical center.**

**PLEASE NOTE:** If you are female patient, please do not urinate prior to coming in for your procedure. You may be asked to provide a urine sample when you arrive at the surgical center.